

Neurotoxicity of Anaesthetics and Sedatives and Their Influence on Post-Operative Maladaptive Behavioural Disorders in Paediatric Anaesthesia (The Letter)

Z. A. Petříková¹, B. Drobná Sáníová², I. Jób³

¹ Department of Paediatric Anaesthesiology and Intensive Care, Children University Hospital
Banská Bystrica, Ludvik Svoboda Square 4, 974 09 Banská Bystrica, Slovak Republic

² Clinic of Anaesthesiology and Intensive Medicine, Comenius University in Bratislava
Jeesenius Faculty of Medicine in Martin, University Hospital,
2 Kollarova Str., 03601 Martin, Slovak Republic

³ Central Slovak Institute for Cardiovascular Diseases, Department of Anaesthesiology and Intensive Care,
1 Cesta k nemocnici, 97401 Banská Bystrica, Slovak Republic

Нейротоксичность анестетиков и седативных средств и их влияние на послеоперационные дезадаптивные расстройства поведения в педиатрической анестезиологии (Письмо в редакцию)

З. А. Петрикова¹, Б. Дробна Саньова², И. Йоб³

¹ Отделение педиатрической анестезиологии и интенсивной терапии, Детская университетская больница,
Словацкая Республика, 97409, г. Банска Быстрица, пл. Людвика Свободы, д. 4

² Братиславский университет им. Коменского, медицинский факультет им. Ессениуса в Мартине,
Университетская клиника, Клиника анестезиологии и интенсивной медицины,
Словацкая Республика, 03601, г. Мартин, ул. Колларова, д. 2

³ Центральный словацкий институт сердечно-сосудистых заболеваний,
отделение анестезиологии и интенсивной терапии,
Словацкая Республика, 97401, г. Банска Бистрица, Дорога в больницу (Cesta k nemocnici), д. 1

For citation: Z. A. Petříková, B. Drobná Sáníová, I. Jób. Neurotoxicity of Anaesthetics and Sedatives and Their Influence on Post-Operative Maladaptive Behavioural Disorders in Paediatric Anaesthesia (The Letter). *Obshchaya Reanimatologiya* = *General Reanimatology*. 2022; 18 (5): 89–93. <https://doi.org/10.15360/1813-9779-2022-5-89-93> [In Russ. and Engl.]

Summary

Neurotoxicity of anaesthetics have become one of the most discussed problems in paediatric anaesthesiology. The experimental studies on animal models have shown that the anaesthetics used in general anaesthesia should have an influence on neurodegenerative processes, neuroapoptosis and the irregular death of the neuronal cells. Because of this fact, scientists are trying to discover the possibilities of how to minimize the adverse effects of anaesthesia and revise the other alternatives of prevention of anaesthesia-induced maladaptive behavioural disorders.

Key words: *neurotoxicity of anaesthetics; maladaptive behavioural disorders; mechanism of neurotoxicity; post-anaesthesia behavioural changes in children; future of paediatric anaesthesiology*

Conflict of interest. The authors declare no conflict of interest.

Read the full-text English version at www.reanimatology.ru

Introduction

The results of recent studies indicate a neurotoxic effect of commonly used inhaled and intravenous anaesthetics on the developing brain of mammals, which has been long overlooked.

In retrospective and observational studies, data appear to indicate behavioural and neurocognitive abnormalities in children who have undergone general anaesthesia. These disorders occur in up to 50% of

children operated on at the youngest age and are resolved within 1 month after the operation. This phenomenon is observed more often the lower is the age of the child, the more painful is the operation and the more restless is the introduction to anaesthesia.

In critically ill new-born children, especially those with low birth weight, a decrease in IQ, a higher incidence of cerebral palsy and visual and hearing impairments have been reported at a later age. Espe-

Correspondence to:

Zuzana Adamcová Petříková
E-mail: zuzankapetrikova@yahoo.com

Адрес для корреспонденции:

Зузана Адамцова Петрикова
E-mail: zuzankapetrikova@yahoo.com

cially at this age, it is very difficult to separate the consequences of stress, homeostasis disorders, surgery and several days of critical condition from the effect of anaesthetics.

Influence of Anaesthetics and Sedatives on Ontogenesis

The development and growth of the mammalian brain is a complex process starting with neurogenesis, continuing with the differentiation of neurons into different subpopulations, migration of nerve cells to their definitive localisation in the central nervous system (CNS), synaptogenesis, synapse formation and myelination of neuron-axon connections. These processes differ significantly depending on the gestational age and species of the mammal in direct relation to the expected life expectancy of the mammal.

Synaptogenesis begins in humans in the third trimester of pregnancy. Brain growth ends at the age of 2–3 years. During physiological development, CNS neurons are produced in considerable abundance. Their subsequent elimination is crucial for achieving normal brain size as well as its morphology. During ontogenesis, excess neurons are eliminated 50–70% by apoptosis [1].

Laboratory work on *in vitro* cultures demonstrates the dependence of the extent of neurodegenerative changes on the age of the individual, the dose of anaesthetic and the duration of its exposure. Differences in regional distribution and cell-specific deleterious effects of anaesthetics on the developing brain as well as their persistence into adolescence in the dentate gyrus and bulbus nervus olfactorius areas were found. A similar effect is described after long-term administration of drugs (especially benzodiazepines) during treatment in intensive care units.

While the sensitivity of neurons to anaesthesia-induced toxicity corresponds to a maximum in the development of synaptogenesis, the greatest susceptibility of oligodendrocytes during exposure to anaesthesia occurs at the time of myelination. Thus, both CNS components are highly sensitive to apoptotic neurodegeneration [2].

Recent research points to a developmentally determined protective role of microglia during brain development and maturation because, under physiological conditions, the microglia alter the synaptic transmission and plasticity of the brain. Under certain conditions — hypoxia, infection, traumatic brain injury, autoimmune neurodegenerative processes — microglia function is enhanced and can modify synaptic connections and plasticity of memory and learning [3].

Many current research findings point to the neurotoxic effect of commonly used anaesthetics and sedatives on animal models after exposure to

doses of anaesthetics used in paediatric anaesthesiology [4].

The neurotoxicity of anaesthetics in animals, which persists into adulthood, depends on the dose and number of anaesthetics used, the maturity of the developing brain at the time of exposure to the anaesthetic and the presence of other factors, especially inflammatory processes in the body. The combination of all influences increases the sensitivity of the brain to the effect of the anaesthetic [3]. «Pharmaceuticals commonly used in intensive care units and operating theatres, such as isoflurane, benzodiazepines, barbiturates, etomidate, propofol and ketamine, are involved in the development of neurotoxicity in animals» [3].

Although these effects on the human body have not yet been clearly demonstrated, a link between anaesthetic exposure and acquired neurological development disorder in children is evident.

In particular, exposure of children to anaesthetics at an early age causes transient suppression of neurogenesis, ultrastructural abnormalities of synapses and alterations in the development of the signalling and neuroinflammatory neural network, loss of neurons, production of free radicals and alterations of mitochondrial integrity. These side effects on the basis of developing neuronal connections can result in acute neuronal damage as well as long-lasting neurocognitive dysfunctions [3–5]. Cognitive deficits are mainly related to the hippocampal region, where the extent of damage to their neurons is much greater compared to other brain regions.

This provides a possible explanation for the neuroapoptosis following anaesthesia described in this area by most studies, while long-term neurocognitive dysfunctions are described in only a few of them. An increase in the incidence of cell death after anaesthetic exposure does not necessarily lead to a significant reduction in neuronal density in old age. During development, 50–70% of all CNS cells undergo natural cell death, thus maintaining the physiological structure of the CNS. It remains unclear whether anaesthesia accelerates the apoptosis of neurons that are primarily destined for death in physiological degeneration, or damages healthy neurons that are not primarily destined for death. Thus, the fact that cognitive deficit is caused by cell death with consequent loss of neurons, i. e. not only by cell death itself, remains an important finding. At the same time, it remains unclear whether neuroapoptosis is the only cause of cognitive dysfunction [5].

In children, associations have been found between long-term exposure to anaesthetics and sedatives, especially GABAergic, and subsequently lower levels of neurological development until the age of 12–48 months.

A special group among these patients are children with congenital heart and vascular diseases, whose survival has increased by up to 90% compared to the past due to neonatal surgery. In 30–50% of these children, intelligence disorders, major or minor motor dysfunctions, and receptive and expressive speech disorders occur after undergoing cardiac surgery. Memory disorders, speech disorders, counting disorders and visual-motor coordination are observed at the age of school entry and integration of these children.

The Mechanism of Anaesthesia-Induced Neurotoxicity

The effect of most anaesthetics results from their action as NMDA receptor antagonists and/or GABA A agonists. Anaesthesia-induced neurotoxicity is mediated through the mitochondrial apoptotic cascade (the internal part of the anaesthetic-induced apoptotic cascade), which activates the neurotropic cascade and subsequently the cascade leading to the destruction of these receptors. Thus, commonly used anaesthetics cause extensive apoptotic neurodegeneration in various parts of the brain during its development.

The main role of every mitochondrion in the cell is the production of energy through oxidative phosphorylation. However, mitochondria also have many regulatory functions that are important for further survival as well as cell death, as exemplified by the intrinsic apoptotic cascade, which leads to organised and controlled cell death. Activation of this cascade is caused by the release of cytochrome c from the mitochondria into the cytosol. Cytochrome c forms apoptosomes followed by activation of apoptotic protease factor (APAF-1) to form deoxyadenosine triphosphate (dATP) and adenosine triphosphate (ATP). Binding of the apoptosome complex activates procaspase-9. Activated caspase-9 activates caspase-3, resulting in deoxyribonucleic acid (DNA) fragmentation and cell death.

The physiological function of cytochrome c is the transfer of electrons between complex III and IV of the respiratory chain during oxidative phosphorylation. However, in binding to cardiolipin, it can also have peroxidase effects, by inducing the oxidation of hydroperoxycardiolipin thereby contributing to the development of pro-apoptotic stimuli. Oxidative stress contributes significantly to its production.

Andropoulos D. B. et al. showed the internal and external pathway of the apoptotic cascade, as well as the antiapoptotic effects of dexmedetomidine and erythropoietin [2].

Many inhaled (isoflurane, sevoflurane) and intravenous (propofol) anaesthetics increase free radical production in the developing brain. Exposure to anaesthetics, even under conditions of normox-

emia, increases the production of free oxygen and nitrogen radicals in developing neurons, hippocampus, subiculum, and thalamus. Oxidative stress caused by anaesthetics leads to peroxidation of membrane lipids, damage to mitochondria and their impaired integrity.

Binding of apoptosis-induced ligands to cell death receptors activates the external part of the anaesthetic-induced apoptotic cascade. The main pro-apoptotic ligands are: tumour necrosis factor alpha (TNF), Fas and TNF-related apoptosis-inducing ligands (TRAILs). Activation of cellular death receptors by the adoption of the intracellular Fas-associated death domain (FADD) leads to the internalisation of procaspase-8, which results not only in its activation but also in the activation of caspase-3. Exposure of isoflurane together with nitric oxide and simultaneously midazolam up-regulated Fas receptors and activated caspase-8 in the parietal and occipital cortex in an experiment in 7-day-old laboratory rats. While the internal pathway of the apoptotic part of the cascade was activated as early as 2 hours after exposure to this combination of anaesthetics. The time difference was due to the dependence on Fas protein expression and upregulation.

NMDA antagonists and GABA receptor agonists cause neuronal cell death by activating the mitochondrial portion of apoptosis.

Neurotrophins belong to growth factors that determine the survival and differentiation of neurons and the plasticity of synapses. Already in 7-day-old laboratory rats exposed to a 6-hour mixture of isoflurane, nitric oxide and midazolam, a decrease in the activity of their main representative BDNF (brain-derived neurotrophic factor) in the developing thalamus was demonstrated.

There are suspicions that opioids may induce apoptosis of developing neurons. For example, infusion of continuous fentanyl results in an increase in caspase-3 levels in specific brain sections of 5-day-old pigs compared to unexposed individuals. Other studies describe the determining effect of morphine on the developing cortex and amygdala. The area of the hippocampus remains surprisingly spared from these influences. The neurotoxic effects of propofol have also been extensively studied and its induced neuronal apoptosis has been described in both rodents and primates [2].

The immature brain is very susceptible to anaesthesia-induced neuronal apoptosis. The fact remains that despite the fact that some neurons undergo cell death after exposure to anaesthetics, while other neurons survive intact [6] demonstrated that the neurotoxicity of anaesthetics depends more on the age of the neuron than on the age of the organism. They pointed out that postnatal gyrus cells undergoing isoflurane-induced neurodegeneration

are young and relatively immature. They reach the maximum anaesthetic-induced vulnerability at the age of 2 weeks after birth. They have also focused their research on olfactorius bulbous cells, which undergo neurogenesis into adulthood, where they have also demonstrated their susceptibility to anaesthesia-induced apoptosis.

The heterogeneity of susceptibility to neurotoxic effects may vary with age. This fact increases the possibility of confirming the assumption that anaesthesia-induced neurotoxicity depends on the age of the organism at the time of its exposure to the adverse effect. Furthermore, the time of manifestation of the neurotoxic effect may exceed the time of early childhood. Young rhesus macaque which were repeatedly exposed to Sevoflurane developed anxiety behaviour up to 6 months after exposure compared to the unexposed sample. Thus, the study only confirms the effect of general anaesthesia on the development of behavioural disorders with a longer time interval from exposure to anaesthetics in primates [4]. Neurodegenerative and neuroprotective effects of anaesthetics have been described in the study by Andropoulos D. B. et al. [2].

Consequences of Neurotoxicity

The success of current intensive care therapy contributes significantly to the increased survival rate of critically ill patients. However, after overcoming this critical and life-threatening period, a significant proportion of paediatric patients have motor, cognitive, and psychological consequences. «Cognitive deficit after overcoming a critical illness leads to a decrease in IQ, poorer school performance and attention and memory disorders. Risk factors for cognitive impairment in patients after hospitalisation in intensive care units include: artificial lung ventilation, extracorporeal vital signs, traumatic impairment, oncological and neurological diseases, and the use of medication for sedation as well as opiates» [3].

Postnatal neurogenesis is a critical period for the development of learning and memory, which are located mainly in the hippocampus. Exposure to the anaesthetic during peak synaptogenesis significantly reduces synaptic transmission and synapse density, thereby causing inhibition of synaptic transmission.

The development of neurons also depends on the integrity of the CNS and the proper function of the astroglia, so there is a reasonable suspicion that the adverse effects of anaesthesia may also occur through its pathways. The adverse effect of reducing the levels of the excitatory neurotransmitters aspartate and glutamate in the cortex and hippocampus on the modulation of learning and memory is also expected [5].

The focus of current laboratory studies is developing therapeutic and preventive strategies to fight neurotoxicity. Of these, the beneficial effects of dexmedetomidine, the mechanism of upregulation of anti-apoptotic proteins, are evident [2]. Erythropoietin also crosses the blood-brain barrier, stimulates neurogenesis, induces neuronal differentiation, activates neurotropic signalling, and also has anti-apoptotic, antioxidant, and anti-inflammatory properties. Due to the anaesthetic-induced production of inflammatory mediators and the development of oxidative stress described after the use of inhaled anaesthetics (isoflurane), the effects of vitamins (vitamin B3, vitamin D3) and other nutrients are also studied [2].

Conclusion

Anaesthesia-induced neuronal damage should have far-reaching consequences in the future of paediatric patients. Therefore, research into the neurotoxicity of drugs used in patients in paediatric intensive care medicine and anaesthesiology should be aimed at elucidating the absolute and dose-effect of anaesthetic drugs on the developing brain. Understanding the mechanism of toxicity of anaesthetics to the developing nervous system, in particular the mechanism by which anaesthesia impairs brain function for up to one month, would significantly contribute to the prevention and development of therapeutic strategies [5]. Further research will be needed to clarify the stage of hippocampal neurogenesis with which anaesthetics interfere, thereby inducing impaired synapse development and their remodelling as a potential cause of cognitive dysfunction. Research will also focus on neurotoxicity therapy options, preventive methods and protection of the developing brain from neurotoxic effects. Some ongoing preclinical animal studies have already shown positive results. If the neurodegenerative effects of anaesthetics are explicitly demonstrated in the future, it will be important to apply therapy and procedures to maintain safety with minimal risk to the paediatric patient. It is also important to focus on the possibilities of adaptation of paediatric anaesthesiologists and paediatric intensivists to prevention and the possibilities of limiting the potential negative impact of anaesthesia on the neurological development of children.

Given the repeated observations of daily practice, other relevant and informative scoring systems will need to be developed in the near future, with not only value for identifying behavioural changes but also for the speed and quality of recovery from original cognitive function as well as recovery time from potential adverse effects of anaesthesia.

References

1. Moore K.L., Persaud T.V.N. The developing human: clinically oriented embryology. 8th Edition. Saunders. 2007: 536. ISBN: 80-85866-94-3.
2. Andropoulos D.B., Stayer S. Anesthesia for congenital heart disease. 3rd Edition. Wiley. 2015: 816. ISBN: 978-1-118-76825-9.
3. Kamat P.P., Kudchadkar S.R., Simon H.K. Sedative and anesthetic neurotoxicity in infants and young children: not just an operating room concern. *J Pediatr.* 2019; 204: 285–290. DOI: 10.1016/j.jpeds.2018.08.039. PMID: 30270160.
4. O'Leary J.D., Warner D.O. What do recent human studies tell us about the association between anaesthesia in young children and neurodevelopmental outcomes? *Br J Anaesth.* 2017; 119 (3): 458–64. DOI: 10.1093/bja/aex141. PMID: 28969310.
5. Deshui Y., Li L., Yuan W. Neonatal anesthetic neurotoxicity: insight into molecular mechanism of long-term neurocognitive deficits. *Biomed Pharmacother.* 2017; 87: 196–199. DOI: 10.1016/j.biopha.2016.12.062. PMID: 28056424.
6. Hofacer R.D., Deng M., Ward C.G., Joseph B., Hughes E.A., Jiang C., Danzer S.C., Loepke A.W. Cell age-specific vulnerability of neurons to anesthetic toxicity. *Ann Neurol.* 2013; 73 (6): 695–704. DOI: 10.1002/ana.23892. PMID: 23526697.

Received 24.01.2022