

The Intrinsic Network Dynamics Related to Abnormal Delta Rhythm in Consciousness Disorders (Short Review)

Calixto Machado^{1*}, Jose J. Sanchez², Beata Drobna Saniova³, Michal Drobny³, Arthur Schiff⁴

¹ Institute of Neurology and Neurosurgery, Department of Clinical Neurophysiology,
29 y D Vedado, 10400 La Habana, Cuba

² Better Call Agency,
1545 NW 15th Street Road, Miami, FL 33125, USA

³ Clinic of Anaesthesiology and Intensive Medicine, Comenius University in Bratislava,
Jessenius Faculty of Medicine and University Hospital in Martin,
2 Kollarova Str., 03659 Martin, Slovak Republic

⁴ Northside Hospital Gwinnett,
500 Medical Center Blvd, Suite 350, Lawrenceville, GA 30046, USA

For citation: Calixto Machado, Jose J. Sanchez, Beata Drobna Saniova, Michal Drobny, Arthur Schiff. The Intrinsic Network Dynamics Related to Abnormal Delta Rhythm in Consciousness Disorders (Short Review). *Obshchaya Reanimatologiya=General Reanimatology*. 2025; 21 (5): 73–76. <https://doi.org/10.15360/1813-9779-2025-5-2584> [In Engl.]

***Correspondence to:** Calixto Machado, cmachado180652@gmail.com

Summary

Aim: to explore the pathophysiological mechanisms and clinical significance of delta rhythms (≤ 4 Hz) in disorders of consciousness (DOC), including coma, unresponsive wakefulness syndrome (UWS), and minimally conscious state (MCS), as biomarkers for diagnosis, prognosis, and therapeutic targeting.

Materials and Methods. A narrative review was conducted, focusing on experimental and clinical findings related to delta rhythm generation and modulation in the disorder of consciousness (DOC). Emphasis was placed on thalamo-cortical interactions, cortical inhibition, neuromodulatory deficits, and the role of glial cells, neuroinflammation, and metabolic disruptions. Quantitative EEG analysis and advanced neuroimaging were highlighted as key tools for assessing delta activity.

Results. Delta rhythms were found to dominate EEGs across DOC states, with high-amplitude global activity in coma and low-amplitude activity in UWS, indicating cortical suppression and thalamocortical disconnection, respectively. In MCS, reduced delta power and improved connectivity correlated with intermittent purposeful behavior. Therapeutic interventions, including TMS, tACS, and pharmacological agents, showed potential for modulating delta rhythms. Additionally, stochastic resonance emerged as a novel mechanism to stabilize neural networks through noise.

Conclusion. Delta rhythms serve as crucial biomarkers in DOC, offering diagnostic, prognostic, and therapeutic value. Multimodal approaches that integrate EEG, neuroimaging, and mechanistic studies are essential for deepening understanding and improving clinical outcomes in DOC management.

Keywords: Delta rhythm; EEG; disorders of consciousness; neural networks; delta-alpha rhythms; neuroimaging

Conflict of interest. The authors declare no conflict of interest.

Внутренняя динамика сети, связанная с аномальным дельта-ритмом при нарушениях сознания (краткий обзор)

К. Мачадо^{1*}, Х. Санчес², Б. Д. Саниова³, М. Дробны³, А. Шифф⁴

¹ Институт неврологии и нейрохирургии, кафедра клинической нейрофизиологии,
Куба, 10400, г. Ла-Гавана, ул. у D Ведадо, д. 29

² Агентство Better Call,
США, 33125, Флорида, Майами, Северо-западная 15-я ул. Роуд, 1545

³ Клиника анестезиологии и интенсивной медицины, Братиславский университет им. Коменского,
медицинский факультет им. Ессениуса и Университетская клиника в Мартине,
Словацкая Республика, 03601, г. Мартин, ул. Колларова, д. 2

⁴ Больница Нортсайд Гвиннетт,
США, 30046, Джорджия, Лоуренсвилл, бульвар Медицинского центра, д. 500, каб. 350

Резюме

Цель: изучить патофизиологические механизмы и клиническую значимость дельта-ритмов (≤ 4 Гц) при нарушениях сознания, включая кому, синдром безответного бодрствования (вегетативное состояние) и состояние минимального сознания (СМС), в качестве биомаркеров для диагностики, прогноза и целенаправленной терапии.

Материалы и методы. Провели краткий обзор результатов экспериментальных и клинических исследований по изучению генерации и модуляции дельта-ритма при нарушениях сознания. Особое

внимание уделено таламо-кортикальным взаимодействиям, кортикальному торможению, нарушениям нейромодуляции, а также роли глиальных клеток, нейровоспалению и метаболическим нарушениям. Количественный анализ ЭЭГ и усовершенствованную нейровизуализацию выделили в качестве ключевых инструментов для оценки дельта-активности.

Результаты. Установили, что дельта-ритмы доминируют на ЭЭГ при всех состояниях нарушенного сознания, при этом глобальная активность с высокой амплитудой наблюдается при коме, а активность с низкой амплитудой — при вегетативных состояниях, что, соответственно, указывает на подавление активности коры головного мозга и таламокортикальное отключение. При СМС снижение амплитуды волн дельта-ритма и улучшение связи коррелировали с периодическим целенаправленным поведением. Терапевтические вмешательства, включая транскраниальную магнитную стимуляцию, транскраниальную стимуляцию переменным током и фармакологические препараты, демонстрировали потенциал для модуляции дельта-ритмов. Кроме того, стохастический резонанс стал новым механизмом стабилизации нейронных сетей, проходящим сквозь помехи.

Заключение. Дельта-ритмы служат важнейшими биомаркерами состояний нарушенного сознания и представляют диагностическую, прогностическую и терапевтическую ценность. Мультимодальные подходы, объединяющие ЭЭГ, нейровизуализацию и механистические исследования, необходимы для углубления понимания и улучшения клинических результатов при лечении пациентов в состоянии нарушенного сознания.

Ключевые слова: дельта-ритм; ЭЭГ; нарушения сознания; нейронные сети; дельта-альфа-ритмы; нейровизуализация

Конфликт интересов. Авторы заявляют об отсутствии конфликта интересов.

Information about the authors:

Machado Calixto/Каликто Мачадо: <https://orcid.org/0000-0002-0539-5844>

Saniova Drobna Beata/Беата Дробна Саниова: <https://orcid.org/0000-0002-0833-9493>

Drobný Michal/Артур Шифф: <https://orcid.org/0000-0001-931>

Introduction

Disorders of consciousness (DOC), including coma, unresponsive wakefulness syndrome (UWS), and minimally conscious state (MCS), represent complex conditions involving disruptions in neural networks. Delta rhythms are prominent in DOC and indicate underlying pathological states. Understanding the mechanisms and implications of delta activity is essential for improving diagnostics, prognostics, and therapeutic approaches. This review synthesizes recent advances in delta rhythm research, emphasizing their mechanistic underpinnings and clinical relevance [1–4].

Mechanisms of Delta Rhythm Generation

Delta rhythms originate from dynamic interactions within the thalamo-cortical circuitry, influenced by [1, 5–8]:

- **Thalamic contributions:** intrinsic oscillatory properties of thalamocortical neurons, mediated by T-type calcium channel dynamics, play a pivotal role in delta generation. Aberrant oscillations within thalamic relay neurons have been linked to hyperpolarized membrane states, resulting in the pathological emergence of delta rhythms.
- **Cortical dynamics:** cortical neurons contribute to delta activity through synchronized hyperpolarization states, facilitated by GABAergic inhibition. Recent studies have shown that disruptions in interneuron networks can exacerbate the desynchronization seen in DOC.
- **Neuromodulatory inputs:** cholinergic and dopaminergic systems modulate delta rhythms, with deficits in these pathways observed in DOC

patients. Furthermore, noradrenergic and serotonergic systems may indirectly influence delta activity by modulating the arousal network [9–13].

- **Role of glial cells:** emerging evidence suggests that astrocytes and microglia contribute to rhythm generation by modulating synaptic activity and clearing metabolic byproducts. Neuroinflammation, a common feature in DOC, may alter glial function and impact delta rhythm dynamics [14–16].

Delta Rhythms in Coma

Coma, a profound state of unresponsiveness, is often characterized by pronounced delta activity [17–20]. Key features include:

- **Global cortical suppression:** coma patients typically exhibit high-amplitude, generalized delta rhythms on EEG, reflecting widespread cortical and subcortical dysfunction.
- **Thalamic dysfunction:** reduced thalamic input in coma is associated with the dominance of slow-wave activity, disrupting normal thalamo-cortical interactions.
- **Prognostic insights:** while persistent delta activity in coma is indicative of severe brain dysfunction, the gradual re-emergence of faster rhythms may signal recovery potential. Studies have highlighted that early shifts in EEG patterns can predict outcomes in coma patients.

Delta Rhythms in Different DOC States [1, 3, 21]

Distinct delta rhythm profiles differentiate coma, UWS, and MCS:

- **Unresponsive wakefulness syndrome (UWS):** Characterized by widespread, high-amplitude delta

activity, reflecting severe cortical disconnection and reduced thalamo-cortical communication.

- *Minimally conscious state (MCS)*: Lower delta power and greater functional connectivity compared to UWS, indicating partial preservation of cortical networks. This preservation is often associated with sporadic evidence of purposeful behavior.
- *Emerging patterns in recovery*: Patients transitioning from DOC states often exhibit gradual reductions in delta power, accompanied by increased alpha and beta rhythms, indicative of cortical reorganization.

Diagnostic and Prognostic Value [22–25]

Delta rhythms serve as biomarkers for:

- **Diagnosis**: Quantitative EEG measures, such as delta power, coherence, and cross-frequency coupling, aid in differentiating DOC states. Combining delta metrics with machine learning approaches has shown promise in improving diagnostic accuracy.
- **Prognosis**: Changes in delta rhythm characteristics over time correlate with recovery potential, informing clinical decision-making. For instance, a shift from global delta dominance to localized delta activity may indicate partial network restoration.

Therapeutic Implications [1, 20, 26, 27]

1. **Neuromodulation**: Techniques such as transcranial magnetic stimulation (TMS) and transcranial alternating current stimulation (tACS) target delta activity, showing promise in restoring cortical network function. The use of personalized stimulation protocols, based on individual EEG profiles, may enhance therapeutic outcomes.

2. **Pharmacological Approaches**: Agents modulating T-type calcium channels or enhancing cholinergic activity offer potential for delta rhythm normalization. Recent trials involving ampakines and GABA-A receptor modulators have demonstrated preliminary efficacy in altering delta dynamics.

3. **Noise Modulation**: Harnessing the paradoxical stabilizing effects of noise, such as through stochastic resonance, represents an emerging therapeutic avenue. Experimental studies suggest that introducing controlled noise can enhance signal transmission in impaired neural circuits.

Литература

1. Wang J., Deng B., Wang J., Yang Y., Liu C. The abnormal delta rhythm mechanism in disorders of consciousness: intrinsic neuronal properties and network dynamics. *Neurocomputing*. 2025; 620: 129206. DOI: 10.1016/j.neucom.2024.129206.
2. Casarotto S., Hassan G., Rosanova M., Sarasso S., Derchi C. C., Trimarchi P. D., Vignani A., et al. Dissociations between spontaneous electroencephalographic features and the perturbational complexity index in the minimally conscious state. *Eur J Neurosci*. 2024; 59: 934–947. DOI: 10.1111/ejn.16299. PMID: 38440949.
3. Bitar R., Khan U. M., Rosenthal S. Utility and rationale for continuous EEG monitoring: a primer for the general intensivist. *Crit Care*. 2024; 28 (244): 1–23. DOI: 10.1186/s13054-024-04986-0. PMID: 39014421.

Integration with Multimodal Biomarkers [28–32]

Combining delta rhythm analysis with neuroimaging modalities enhances the understanding of DOC pathophysiology. For instance:

- **Functional MRI (fMRI)**: reveals connectivity patterns linked to delta activity, providing insights into network-level disruptions.
- **Diffusion Tensor Imaging (DTI)**: assesses white matter integrity associated with rhythm generation. Studies have shown that reduced fractional anisotropy in thalamic and cortical regions correlates with elevated delta power.
- **PET Imaging**: investigates metabolic abnormalities that may underlie delta rhythm alterations, such as glucose hypometabolism in key brain regions.

Future Directions

Key areas for further exploration include:

- **Mechanistic studies**: elucidating the interplay of glial cells, neuroinflammation, and vascular dynamics in delta rhythm generation. Investigating the role of extracellular potassium accumulation in modulating delta activity may also be fruitful.
- **Longitudinal studies**: examining delta rhythm changes during recovery and their relationship to neural plasticity. Such studies could identify biomarkers predictive of long-term outcomes.
- **Clinical trials**: evaluating the efficacy of delta-modulating interventions in large patient cohorts. Trials should aim to stratify patients based on baseline EEG features to optimize therapeutic targeting.
- **Technological innovations**: developing portable, high-density EEG systems to facilitate real-time monitoring of delta activity in clinical settings.

Conclusion

Delta rhythms offer profound insights into the neural disruptions underlying DOC. Advances in understanding their mechanisms and clinical implications hold promise for improving patient outcomes. Future research should focus on bridging experimental findings with translational applications to harness the diagnostic and therapeutic potential of delta rhythms fully. Integrating delta rhythm analysis with multimodal approaches and personalized therapies will likely define the next frontier in DOC management.

4. Valomon A., Riedner B. A., Jones S. G., Nakamura K. P., Tononi G., Plante D. T., Benca R. M., et al. high-density electroencephalography study reveals abnormal sleep homeostasis in patients with rapid eye movement sleep behavior disorder. *Sci Rep*. 2021; 11: 4758. DOI: 10.1038/s41598-021-83980-w. PMID: 33637812.
5. Naro A., Bramanti A., Leo A., Cacciola A., Manuli A., Bramanti P., Calabro R. S. Shedding new light on disorders of consciousness diagnosis: The dynamic functional connectivity. *Cortex*. 2018; 103: 316–328. DOI: 10.1016/j.cortex.2018.03.029. PMID: 29684751.
6. De Stefano P., Carboni M., Pugin D., Seck M., Vulliemoz S. Brain networks involved in generalized periodic discharges (GPD) in post-anoxic-ischemic encephalopathy. *Resuscitation*. 2020; 155: 143–151. DOI: 10.1016/j.resuscitation.2020.07.030. PMID: 32795598.

7. Llinás R. R. Thalamo-cortical dysrhythmia syndrome: neuropsychiatric features. *An R Acad Nac Med (Madr.)*. 2003; 120 (2): 267–290. PMID: 14619543.
8. Jones B. E. Arousal systems. *Front Biosci*. 2003; 8: s438–451. DOI: 10.2741/1074. PMID: 12700104.
9. Brudzynski S. M., Burgdorf J. S., Moskal J. R. From emotional arousal to executive action. Role of the prefrontal cortex. *Brain Struct Funct*. 2024; 229: 2327–2338. DOI: 10.1007/s00429-024-02837-w. PMID: 39096390.
10. Babiloni C., Del Percio C., Pascarelli M. T., Lizio R., Noce G., Lopez S., Rizzo M., et al. Abnormalities of functional cortical source connectivity of resting-state electroencephalographic alpha rhythms are similar in patients with mild cognitive impairment due to Alzheimer's and Lewy body diseases. *Neurobiol Aging*. 2019; 77: 112–127. DOI: 10.1016/j.neurobiolaging.2019.01.013. PMID: 30797169.
11. Carracedo L. M., Kjeldsen H., Cunningham L., Jenkins A., Schofield I., Cunningham M. O., Davies C. H., et al. A neocortical delta rhythm facilitates reciprocal interlaminar interactions via nested theta rhythms. *J Neurosci*. 2013; 33 (26): 10750–10761. DOI: 10.1523/JNEUROSCI.0735-13.2013. PMID: 23804097.
12. Dimpfel W. Pharmacological modulation of cholinergic brain activity and its reflection in special EEG frequency ranges from various brain areas in the freely moving rat (Tele-Stereo-EEG). *Eur Neuropsychopharmacol*. 2005; 15 (6): 673–682. DOI: 10.1016/j.euroneuro.2005.03.006. PMID: 16009538.
13. Philippu A., Prast H. Importance of histamine in modulatory processes, locomotion and memory. *Behav Brain Res*. 2001; 124 (2): 151–159. DOI: 10.1016/s0166-4328 (01)00226-1. PMID: 11640968.
14. Hao W., Liu S., Liu H., Mu X., Chen K., Xin Q., Zhang X.-D. In vivo neuroelectrophysiological monitoring of atomically precise Au (25) clusters at an ultrahigh injected dose. *ACS Omega*. 2020; 5 (38): 24537–24545. DOI: 10.1021/acsomega.0c03005. PMID: 33015471.
15. Grigorevsky V., Bardakjian B. L. Low-to-high cross-frequency coupling in the electrical rhythms as biomarker for hyperexcitable neuroglial networks of the brain, IEEE. *Trans Biomed Eng*. 2018; 65 (7): 1504–1515. DOI: 10.1109/TBME.2017.2757878. PMID: 28961101.
16. Gavilanes A. W., Gantert M., Strackx E., Zimmermann L. J., Seelgraber S., Vles J. S., Kramer B. W. Increased EEG delta frequency corresponds to chorioamnionitis-related brain injury. *Front Biosci (Schol Ed)*. 2010; 2 (2): 432–438. DOI: 10.2741/s76. PMID: 20036959.
17. Kumar G. P., Panda R., Sharma K., Adarsh A., Annen J., Martial C., Faymonville M. E., et al. Changes in high-order interaction measures of synergy and redundancy during non-ordinary states of consciousness induced by meditation, hypnosis, and auto-induced cognitive trance. *Neuroimage*. 2024; 293: 120623, 1–10. DOI: 10.1016/j.neuroimage.2024.120623. PMID: 38670442.
18. Rayi A., Asuncion R. M. D., Mandalaneni K. Encephalopathic EEG patterns. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025. 2024 Aug 11. PMID: 33232041. Bookshelf ID: NBK564371.
19. Huang Y., Xia X., Meng X., Bai Y., Feng Z. Single session of intermittent theta burst stimulation alters brain activity of patients in vegetative state. *Aging*. 2024; 16 (8): 7119–7130. DOI: 10.18632/aging.205746. PMID: 38643463.
20. Li M., Tang Z., Yu L., Li Y., Ma W., Li J., Li G., Xiong L., et al. The arousal effect of An-Gong-Niu-Huang-Wan on alcoholic-induced coma rats: a research based on EEG. *J Ethnopharmacol*. 2024; 328: 117974. DOI: 10.1016/j.jep.2024.117974. PMID: 38467317.
21. Espinosa E. P., Zang D., Buccellato A., Qi Z., Wu X., Abbasi S., Catal Y., et al. Spectral peak analysis and intrinsic neural timescales as markers for the state of consciousness. *Neuroimage Clin*. 2024; 44: 103698. DOI: 10.1016/j.nicl.2024.103698. PMID: 39509990.
22. Li C., Wang Y., Li W., Yang Y., Xia X. Measure functional network and cortical excitability in post-anoxic patients with unresponsive wakefulness syndrome diagnosed by behavioral scales. *Front Neurosci*. 2022; 16: 1071594. DOI: 10.3389/fnins.2022.1071594. PMID: 36711155.
23. Bai Y., Xia X., Wang Y., He J., X. Li X. Electroencephalography quadratic phase self-coupling correlates with consciousness states and restoration in patients with disorders of consciousness. *Clin Neurophysiol*. 2019; 130: 1235–1242. DOI: 10.1016/j.clinph.2019.04.710. PMID: 31163368.
24. Stefan S., B. Schorr B., Lopez-Rolon A., Kolassa I. T., Shock J. P., Rosenfelder M., Heck S., et al. Consciousness indexing and outcome prediction with resting-state EEG in severe disorders of consciousness. *Brain Topogr*. 2018; 31 (5): 848–862. DOI: 10.1007/s10548-018-0643-x. PMID: 29666960.
25. Bagnato S., Boccagni C., Prestandrea C., Fingelkurts A. A., Fingelkurts A. A., Galardi G. Changes in standard electroencephalograms parallel consciousness improvements in patients with unresponsive wakefulness syndrome. *Arch Phys Med Rehabil*. 2017; 98 (4): 665–672. DOI: 10.1016/j.apmr.2016.09.132. PMID: 27794486.
26. Anzolin A., Das P., Garcia R. G., Chen A., Grahl A., Ellis S., Purdon P., et al. Delta power during sleep is modulated by EEG-gated auricular vagal afferent nerve stimulation (EAVANS). *Annu Int Conf IEEE Eng Med Biol Soc*. 2023: 1–4. DOI: 10.1109/EMBC40787.2023.10340971. PMID: 38082663.
27. Landau J., Baulac M., Durand G., A. de Billy A., Philippon J. Impairment of consciousness induced by valproate treatment following neurosurgical operation. *Acta Neurochir (Wien)*. 1993; 125 (1–4): 92–96. DOI: 10.1007/BF01401834. PMID: 7993403.
28. Bonfiglio L., Olcese U., Rossi B., Frisoli A., Arrighi P., Greco G., Carozzo S., et al. Cortical source of blink-related delta oscillations and their correlation with levels of consciousness. *Hum Brain Mapp*. 2013; 34 (9): 2178–2189. DOI: 10.1002/hbm.22056. PMID: 22431380.
29. Duda A. T., Clarke A. R., Barry R. J., De Blasio F. M. Mindfulness meditation is associated with global EEG spectral changes in theta, alpha, and beta amplitudes. *Int J Psychophysiol*. 2024; 206: 112465. DOI: 10.1016/j.ijpsycho.2024.112465. PMID: 39557128.
30. Babiloni C., Lopez S., Noce G., Ferri R., Panerai S., Catania V., Soricelli A., et al. Relationship between default mode network and resting-state electroencephalographic alpha rhythms in cognitively unimpaired seniors and patients with dementia due to Alzheimer's disease. *Cereb Cortex*. 2023; 33 (20): 10514–10527. DOI: 10.1093/cercor/bhad300. PMID: 37615301.
31. Belov D., Fesenko Z., Efimov A., Lakstygala A., Efimova E. Different sensitivity to anesthesia according to ECoG data in dopamine transporter knockout and heterozygous rats. *Neurosci Lett*. 2022; 788: 136839. DOI: 10.1016/j.neulet.2022.136839. PMID: 35964824.
32. Jombik P., Drobny M., Saniova B., Fischer M., Kaderjakova P., Lajciakova M., Bakosova E., et al. Some quantitative EEG features in default mode resting state network under general anaesthesia. *Neuro Endocrinol Lett*. 2017; 38 (4): 261–268. PMID: 28871712.

Поступила 24.01.2025

Принята 17.09.2025

Опубликовано онлайн 02.10.2025